

Generalized local cohomology and the Intersection Theorem

Mohammad T. Dibaei ^{ab} and Siamak Yassemi^{ca}

(a) *Institute for Studies in Theoretical Physics and Mathematics, P.O. Box 19395-5746, Tehran, Iran*

(b) *Department of Mathematics, Teacher Training University, Tehran, Iran*

(c) *Department of Mathematics, University of Tehran, Tehran, Iran*

Abstract

Let R be commutative Noetherian ring and let $\mathfrak{a}$ be an ideal of R . For complexes X and Y of R -modules we investigate the invariant $\inf \mathbf{R}\Gamma_{\mathfrak{a}}(\mathbf{R}\mathrm{Hom}_R(X, Y))$ in certain cases. It is shown that, for bounded complexes X and Y with finite homology, $\dim Y \leq \dim \mathbf{R}\mathrm{Hom}_R(X, Y) \leq \mathrm{proj.dim} X + \dim(X \otimes_R^{\mathbf{L}} Y) + \sup X$ which strengthen the Intersection Theorem. Here $\inf X$ and $\sup X$ denote the homological infimum, and supremum of the complex X , respectively.

0. Introduction

Cohomological dimension of a module M over a finite dimensional Noetherian ring R with respect to an ideal $\mathfrak{a}$ is denoted by $\mathrm{cd}(\mathfrak{a}, M) = \sup\{i \in \mathbb{Z} | H_{\mathfrak{a}}^i(M) \neq 0\}$ and has been studied by Faltings [Fa] and Huneke–Lyubeznik [HL], where $H_{\mathfrak{a}}^i(M)$ is the i -th right derived functor of the section functor $\Gamma_{\mathfrak{a}}(M)$. The authors, in [DY], introduced the notion of cohomological dimension, $\mathrm{cd}(\mathfrak{a}, X)$, of a bounded to the left complex X with respect to an ideal $\mathfrak{a}$ to be $\sup\{\mathrm{cd}(\mathfrak{a}, H_{\ell}(X)) - \ell | \ell \in \mathbb{Z}\}$. It is shown that $-\inf \mathbf{R}\Gamma_{\mathfrak{a}}(X) \leq \mathrm{cd}(\mathfrak{a}, X)$ and equality holds whenever X has finite homology modules.

Let M be a finite R -module with finite projective dimension and let N be a finite R -module. A consequence of the New Intersection Theorem from [PS] is the dimension inequality:

2000 *Mathematics subject classification.* 13D45, 13D22, 13D25, 13D05.

Key words and phrases. Intersection Theorem, cohomological dimension, complexes of modules.

The research of the first author was supported by a grant from IPM (No. 82130014).

The second author was supported by a grant from University of Tehran (No. 511/3/702).

Email addresses: dibaeimt@ipm.ir and yassemi@ipm.ir

$\dim N \leq \text{proj.dim } M + \dim (M \otimes_R N)$, cf. [R; 8.4.4]. Actually the following inequalities hold (see [Y1; Remark 3.8]):

$$(T1) \quad \dim N \leq \dim \mathbf{RHom}_R(M, N) \leq \dim M \otimes_R N + \text{proj.dim } M.$$

In particular, if we replace N by R then we have

$$(T2) \quad \dim R \leq \dim \mathbf{RHom}_R(M, R) \leq \dim M + \text{proj.dim } M.$$

Since the notions “dimension” and “cohomological dimension with respect to an ideal” are closed in some sense, it is natural to ask about the following question: “Do we have the inequality $\text{cd}(\mathfrak{a}, \mathbf{RHom}_R(M, N)) \leq \text{cd}(\mathfrak{a}, M \otimes_R N) + \text{proj.dim } M$ for finite R -modules M and N with $\text{proj.dim } M < \infty$?” In this paper we give a positive answer to this question.

In 1967 Auslander [A] introduced a new invariant for finite modules, a relative homological dimension, called the Gorenstein dimension and denoted by $\text{Gdim}(-)$. It is well-known that for all finite R -modules M there is an inequality $\text{Gdim } M \leq \text{proj.dim } M$ and equality holds if $\text{proj.dim } M < \infty$. The Gorenstein dimension is extended for complexes in [Y2] (see also [C]).

After introductory section 1, we consider complexes of finite homologies and show in section 2 that if X and Y are bounded complexes, then

$$\dim \mathbf{RHom}_R(X, Y) \leq \dim R + \sup X - \inf Y$$

provided $\text{proj.dim } X$ or $\text{inj.dim } Y$ is finite, or, $\text{Gdim } X$ and $\text{proj.dim } Y$ are finite. This result, which is a generalization of [HZ; Theorem 3.2], implies the equality $\dim \mathbf{RHom}_R(M, R) = \dim R$, provided $\text{proj.dim } M < \infty$, which forces the left inequality of (T2) to be an equality. This result motivates us to give some characterization properties for a local ring $(R, \mathfrak{m}, k)$. For example, $(R, \mathfrak{m}, k)$ is Gorenstein (resp. Cohen–Macaulay) ring if and only if $\dim \mathbf{RHom}_R(X, R)$ is finite (resp. $\dim \mathbf{RHom}_R(X, L)$ is finite for some non-zero finite R -module L) for all bounded complexes X with finite homologies (see 2.3 and 2.4).

We next examine the Intersection Theorem and show that

$$\dim Y + \inf X \leq \dim \mathbf{RHom}_R(X, Y) \leq \dim (X \otimes_R^{\mathbf{L}} Y) + \text{proj.dim } X + \sup X$$

for all bounded complexes X, Y with finite homologies and $\text{proj.dim } X < \infty$ (see Theorem 2.6). This result generalizes (T1), which in particular implies

$$\dim Y \leq \dim \mathbf{RHom}_R(M, Y) \leq \dim M \otimes_R^{\mathbf{L}} Y + \text{proj.dim } M$$

for all bounded complexes Y and all finite R -module M with finite projective dimension (see Corollary 2.7).

In section 3, we mainly investigate the cohomological dimension of $\mathbf{R}\mathrm{Hom}_R(X, Y)$ with respect to an ideal $\mathfrak{a}$ of R , where X, Y are appropriate R -complexes with $\mathrm{proj.dim} X < \infty$. Our main result is as follows:

If X, Y are bounded complexes with finite homologies then

$$\begin{aligned} -\inf \mathbf{R}\Gamma_{\mathfrak{a}}(Y) + \inf X &\leq -\inf \mathbf{R}\Gamma_{\mathfrak{a}}(\mathbf{R}\mathrm{Hom}_R(X, Y)) \\ &\leq -\inf \mathbf{R}\Gamma_{\mathfrak{a}}(X \otimes_R^{\mathbf{L}} Y) + \mathrm{proj.dim} X + \sup X, \end{aligned}$$

where the first inequality holds under the extra condition $\mathrm{Supp} Y \subseteq \mathrm{Supp} X$ (see Theorems 3.2 and 3.3).

Our results in section 3 show the connection between cohomological dimension and the Intersection Theorem.

Throughout R is a commutative Noetherian ring and $\mathfrak{a}$ is an ideal of R .

1. Notations

An R -complex X is a sequence of R -modules X_{ℓ} and R -linear maps $\partial_{\ell}^X, \ell \in \mathbb{Z}$,

$$X = \cdots \longrightarrow X_{\ell+1} \xrightarrow{\partial_{\ell+1}^X} X_{\ell} \xrightarrow{\partial_{\ell}^X} X_{\ell-1} \longrightarrow \cdots .$$

The module X_{ℓ} is called the module in degree ℓ , and the map $\partial_{\ell}^X : X_{\ell} \longrightarrow X_{\ell-1}$ is the ℓ -th differential, and $\partial_{\ell}^X \partial_{\ell+1}^X = 0$ for all $\ell \in \mathbb{Z}$. An R -module M is thought of as the complex $M = 0 \longrightarrow M \longrightarrow 0$.

The *supremum* and *infimum* of X are defined by

$$\begin{aligned} \sup X &= \sup \{\ell \in \mathbb{Z} | H_{\ell}(X) \neq 0\} \\ \inf X &= \inf \{\ell \in \mathbb{Z} | H_{\ell}(X) \neq 0\}, \end{aligned}$$

denote $\sup X = -\infty$ and $\inf X = \infty$ if $H_{\ell}(X) = 0$ for all ℓ ; in this case X is called homologically trivial.

A morphism $\alpha : X \longrightarrow Y$ is said to be a quasi-isomorphism, and denoted by $X \simeq Y$, if the induced morphism $H(\alpha) : H(X) \longrightarrow H(Y)$ is an isomorphism. Support of X is defined by $\mathrm{Supp} X = \{\mathfrak{p} \in \mathrm{Spec} R | X_{\mathfrak{p}} \text{ is not homologically trivial}\}$.

The *derived category* $\mathcal{D}(R)$ is the category of R -complexes localized at the class of all quasi-isomorphisms. The full subcategories $\mathcal{D}_+(R)$, $\mathcal{D}_-(R)$, and $\mathcal{D}_b(R)$ of $\mathcal{D}(R)$ consist of complexes X with $H_\ell(X) = 0$ for, respectively, $\ell \ll 0$, $\ell \gg 0$, and $|\ell| \gg 0$. By $\mathcal{D}^f(R)$ we mean the full subcategory of $\mathcal{D}(R)$ consisting of complexes X with $H_\ell(X)$ is a finite R -module for all ℓ .

The left derived functor of the tensor product functor of R -complexes is denoted by $-\otimes_R^{\mathbf{L}}-$, and $\mathbf{R}\mathrm{Hom}_R(-, -)$ denotes the right derived functor of the homomorphism functor of complexes. We need the next two inequalities for $X, Y \in \mathcal{D}_+(R)$ and $Z \in \mathcal{D}(R)$.

$$(1.1) \quad \inf(X \otimes_R^{\mathbf{L}} Y) \geq \inf X + \inf Y \quad \text{and}$$

$$(1.2) \quad \sup(\mathbf{R}\mathrm{Hom}_R(X, Z)) \leq \sup Z - \inf X.$$

For a complex X , the dimension of X is defined by Foxby in [Fo] as follows:

$$(1.3) \quad \dim_R X = \sup\{\dim R/\mathfrak{p} - \inf X_{\mathfrak{p}} | \mathfrak{p} \in \mathrm{Spec} R\}.$$

It is shown, in [Fo; 16.9], that:

$$\dim X = \sup\{\dim_R H_\ell(X) - \ell | \ell \in \mathbb{Z}\}.$$

Therefore it is natural to give the following definition, cf. [DY; 2.1]: For a complex $X \in \mathcal{D}_+(R)$, the $\mathfrak{a}$ -cohomological dimension of X is defined by

$$\mathrm{cd}(\mathfrak{a}, X) = \sup\{\mathrm{cd}(\mathfrak{a}, H_\ell(X)) - \ell | \ell \in \mathbb{Z}\}.$$

Note that for an R -module M , this notion agrees with the classical one.

A complex $X \in \mathcal{D}_b(R)$ is said to be of finite *projective* (resp. *injective*) *dimension* if $X \simeq U$, where $U \in \mathcal{D}_b(R)$ is a complex of projective (resp. injective) modules.

The full subcategories of $\mathcal{D}_b(R)$ consisting of complexes of finite projective (resp. injective) dimension are denoted by $\mathcal{P}(R)$ (resp. $\mathcal{I}(R)$). If X belongs to $\mathcal{D}_b(R)$, then the following inequalities hold when $P \in \mathcal{P}(R)$, $I \in \mathcal{I}(R)$, cf. [Fo; 8.9 and 8.13].

$$\inf(\mathbf{R}\mathrm{Hom}_R(P, X)) \geq \inf X - \mathrm{proj.dim}_R P;$$

$$\inf(\mathbf{R}\mathrm{Hom}_R(X, I)) \geq -\sup X - \mathrm{inj.dim}_R I.$$

2. Dimensions and Intersection Theorem

In this section we study dimension of the R -complex $\mathbf{R}\mathrm{Hom}_R(X, Y)$ for appropriate complexes X and Y . For motivation, let us assume M be a finite R -module of finite projective dimension. In [HZ] Herzog and Zamani, by using Buchsbaum–Eisenbud criterion, show that $\dim \mathrm{Ext}_R^i(M, R) \leq \dim R - i$ for all $i \in \mathbb{Z}$, which is equivalent to say $\dim \mathbf{R}\mathrm{Hom}_R(M, R) \leq \dim R$. Here, we first bring a generalization of this result without using of the mentioned criterion; and then we find some characterizations of complexes in $\mathcal{D}_b^f(R)$ to have finite projective (or injective) dimensions.

In [AB], Auslander and Bridger have introduced the notion of Gorenstein dimension of a finite R -module M as $\mathrm{Gdim} M = \sup\{i \in \mathbb{Z} | \mathrm{Ext}_R^i(M, R) \neq 0\}$, and generalized in [Y2] for reflexive complexes. A complex $X \in \mathcal{D}(R)$ is called reflexive if $X \in \mathcal{D}_b^f(R)$, $\mathbf{R}\mathrm{Hom}_R(X, R) \in \mathcal{D}_b^f(R)$ and the natural map $X \rightarrow \mathbf{R}\mathrm{Hom}_R(\mathbf{R}\mathrm{Hom}_R(X, R), R)$ is a quasi-isomorphism. For a reflexive complex X , Gorenstein dimension of X is defined by $\mathrm{Gdim} X = -\inf \mathbf{R}\mathrm{Hom}(X, R)$, cf. [Y2; Definition 2.8]. Note that $\mathrm{Gdim} X \leq \mathrm{proj.dim} X$, and equality holds if $\mathrm{proj.dim} X < \infty$.

Theorem 2.1. Let $X, Y \in \mathcal{D}_b^f(R)$. Then

$$\dim \mathbf{R}\mathrm{Hom}_R(X, Y) \leq \dim R + \sup X - \inf Y$$

provided one of the following conditions holds:

- (i) $\mathrm{proj.dim} X$ is finite,
- (ii) $\mathrm{inj.dim} Y$ is finite,
- (iii) $\mathrm{Gdim} X$ and $\mathrm{proj.dim} Y$ are finite.

Proof. We may assume that $\mathbf{R}\mathrm{Hom}_R(X, Y)$ is a homologically non-trivial complex. Choose an integer ℓ with $H_\ell(\mathbf{R}\mathrm{Hom}_R(X, Y)) \neq 0$. Then $\mathrm{grade} H_\ell(\mathbf{R}\mathrm{Hom}_R(X, Y)) = \mathrm{depth} R_{\mathfrak{p}}$ for some $\mathfrak{p} \in \mathrm{Supp} H_\ell(\mathbf{R}\mathrm{Hom}_R(X, Y))$.

- (i) Assume $\mathrm{proj.dim} X < \infty$. As $H_\ell(\mathbf{R}\mathrm{Hom}_{R_{\mathfrak{p}}}(X_{\mathfrak{p}}, Y_{\mathfrak{p}})) \neq 0$, we have

$$-\ell \leq -\inf \mathbf{R}\mathrm{Hom}_{R_{\mathfrak{p}}}(X_{\mathfrak{p}}, Y_{\mathfrak{p}}) \leq \mathrm{proj.dim} X_{\mathfrak{p}} - \inf Y_{\mathfrak{p}}.$$

Set $H_\ell(\mathbf{R}\mathrm{Hom}_R(X, Y)) = T$. By using Auslander–Buchsbaum formula, $\mathrm{proj.dim} X_{\mathfrak{p}} = \mathrm{depth} R_{\mathfrak{p}} - \mathrm{depth}_{R_{\mathfrak{p}}} X_{\mathfrak{p}}$, and so the following computation hold:

$$\begin{aligned} \dim T - \ell &\leq \dim T + \mathrm{proj.dim} X_{\mathfrak{p}} - \inf Y_{\mathfrak{p}} \\ &= \dim T + \mathrm{depth} R_{\mathfrak{p}} - \mathrm{depth}_{R_{\mathfrak{p}}} X_{\mathfrak{p}} - \inf Y_{\mathfrak{p}} \\ &= \dim T + \mathrm{grade} T - \mathrm{depth}_{R_{\mathfrak{p}}} X_{\mathfrak{p}} - \inf Y_{\mathfrak{p}} \\ &\leq \dim R - \mathrm{depth}_{R_{\mathfrak{p}}} X_{\mathfrak{p}} - \inf Y_{\mathfrak{p}}. \end{aligned}$$

By [Fo; 12.6] we have $-\mathrm{depth}_{R_{\mathfrak{p}}} X_{\mathfrak{p}} \leq \sup X_{\mathfrak{p}} \leq \sup X$. Now by using the fact that $\inf Y \leq \inf Y_{\mathfrak{p}}$, we get the assertion.

(ii) Assume $\mathrm{inj.dim} Y < \infty$. We have

$$-\ell \leq -\inf \mathbf{R}\mathrm{Hom}_{R_{\mathfrak{p}}}(X_{\mathfrak{p}}, Y_{\mathfrak{p}}) \leq \mathrm{inj.dim} Y_{\mathfrak{p}} + \sup X_{\mathfrak{p}}.$$

Again let $H_\ell(\mathbf{R}\mathrm{Hom}_R(X, Y)) = T$. Now, using [Fo; 13.23 (I')] implies that

$$\begin{aligned} \dim T - \ell &\leq \dim T + \mathrm{inj.dim} Y_{\mathfrak{p}} + \sup X_{\mathfrak{p}} \\ &= \dim T + \mathrm{depth} R_{\mathfrak{p}} - \inf Y_{\mathfrak{p}} + \sup X_{\mathfrak{p}} \\ &\leq \dim T + \mathrm{grade} T - \inf Y + \sup X \\ &\leq \dim R - \inf Y + \sup X. \end{aligned}$$

Thus the assertion holds.

(iii) Assume $\mathrm{Gdim} X < \infty$ and $\mathrm{proj.dim} Y < \infty$. By [C; Proposition 2.4.1 and Theorem 2.3.13], we have

$$\begin{aligned} -\ell \leq -\inf \mathbf{R}\mathrm{Hom}_{R_{\mathfrak{p}}}(X_{\mathfrak{p}}, Y_{\mathfrak{p}}) &\leq \mathrm{Gdim} X_{\mathfrak{p}} - \inf Y_{\mathfrak{p}} \\ &= \mathrm{depth} R_{\mathfrak{p}} - \mathrm{depth}_{R_{\mathfrak{p}}} X_{\mathfrak{p}} - \inf Y_{\mathfrak{p}}. \end{aligned}$$

Now the assertion follows with the same argument as in (i). $\square$

The next Corollary shows that the equality holds for the left inequality in (T2).

Corollary 2.2 Assume that M is a non-zero finite R -module with $\mathrm{proj.dim} M < \infty$. Then $\dim \mathbf{R}\mathrm{Hom}_R(M, R) = \dim R$.

Proof. Use Theorem 2.1(i) and (T2). $\square$

In [AB], Auslander and Bridger show that the ring R is Gorenstein if and only if all finite R -modules have finite Gorenstein dimension. The following characterization of Gorenstein rings is parallel to the Auslander–Bridger characterization.

Corollary 2.3 Let $(R, \mathfrak{m}, k)$ be a local ring. Then the following statements are equivalent.

- (i) R is a Gorenstein ring;
- (ii) $\dim \mathbf{R}\mathrm{Hom}_R(X, R) < \infty$ for all $X \in \mathcal{D}_b^f(R)$;
- (iii) $\dim \mathbf{R}\mathrm{Hom}_R(M, R) < \infty$ for all finite R -module M ;
- (iv) $\dim \mathbf{R}\mathrm{Hom}_R(k, R) < \infty$.

Proof. (i) $\Rightarrow$ (ii) is clear from Theorem 2.1. (ii) $\Rightarrow$ (iii) and (iii) $\Rightarrow$ (iv) are trivial. For (iv) $\Rightarrow$ (i), note that

$$\sup\{\ell | \mathrm{Ext}_R^\ell(k, R) \neq 0\} = \dim \mathbf{R}\mathrm{Hom}_R(k, R) < \infty,$$

so R is a Gorenstein ring. $\square$

The Bass characterization of Cohen–Macaulay rings says that R is Cohen–Macaulay if and only if there exists a finite R -module with finite injective dimension, cf. [BH; Page 375]. The following characterization of Cohen–Macaulay rings is parallel to the Bass characterization.

Corollary 2.4 Let $(R, \mathfrak{m}, k)$ be a local ring. Then the following statements are equivalent.

- (i) R is a Cohen–Macaulay ring;
- (ii) There exists a non-zero finite R -module L such that $\dim \mathbf{R}\mathrm{Hom}_R(X, L) < \infty$ for all $X \in \mathcal{D}_b^f(R)$;
- (iii) There exists a non-zero finite R -module L such that $\dim \mathbf{R}\mathrm{Hom}_R(M, L) < \infty$ for all finite R -module M ;
- (iv) There exists a non-zero finite R -module L such that $\dim \mathbf{R}\mathrm{Hom}_R(k, L) < \infty$.

Proof. Note that Cohen–Macaulayness of R is equivalent to the existence of a non-zero finite R -module L of finite injective dimension, and use Theorem 2.1. $\square$

Let $(R, \mathfrak{m})$ be local ring. Let M and N be finite R -modules and M has finite projective dimension. The New Intersection Theorem of Peskine and Szpiro [PS], Hochster [H], and P.

Roberts [R] yields an inequality

$$\dim N \leq \dim (M \otimes_R N) + \text{proj.dim } M.$$

Actually by (T1) we have the following inequalities,

$$\dim N \leq \dim \mathbf{R}\text{Hom}_R(M, N) \leq \dim (M \otimes_R N) + \text{proj.dim } M.$$

On the other hand in [Fo] Foxby generalized the New Intersection Theorem for complexes of modules that is

$$\dim Y \leq \dim (X \otimes_R^{\mathbf{L}} Y) + \text{proj.dim } X$$

for any $X, Y \in \mathcal{D}_b^f(R)$ with $\text{proj.dim } X < \infty$. Now we generalize (T1) for complexes. To do this we need the following Theorem.

Theorem 2.5[Fo; 9.6 and 11.27(d)] Assume that $X \in \mathcal{D}_+^f(R) \cap \mathcal{P}(R)$ and $Y \in \mathcal{D}_b(R)$. Then the following hold.

- (a) $\mathbf{R}\text{Hom}_R(X, Y) \simeq \mathbf{R}\text{Hom}_R(X, R) \otimes_R^{\mathbf{L}} Y$
- (b) If R is local then $\inf \mathbf{R}\text{Hom}_R(X, R) = -\text{proj.dim } X$.

Theorem 2.6 If $X, Y \in \mathcal{D}(R)$ such that $\mathbf{R}\text{Hom}_R(X, Y)$ is not homologically trivial and that $\text{proj.dim } X < \infty$, then the following statements hold:

- (i) If $X \in \mathcal{D}_+^f(R)$ and $Y \in \mathcal{D}_b(R)$, then

$$\dim \mathbf{R}\text{Hom}_R(X, Y) \leq \dim Y + \text{proj.dim } X.$$

- (ii) If $X, Y \in \mathcal{D}_b^f(R)$, then

$$\dim Y + \inf X \leq \dim \mathbf{R}\text{Hom}_R(X, Y) \leq \dim (X \otimes_R^{\mathbf{L}} Y) + \text{proj.dim } X + \sup X.$$

Proof. Choose a prime ideal $\mathfrak{p} \in \text{Supp } X \cap \text{Supp } Y$ such that

$$\begin{aligned}
\dim \mathbf{R}\mathrm{Hom}_R(X, Y) &= \dim R/\mathfrak{p} - \inf \mathbf{R}\mathrm{Hom}_R(X, Y)_{\mathfrak{p}} \\
&= \dim R/\mathfrak{p} - \inf \mathbf{R}\mathrm{Hom}_{R_{\mathfrak{p}}}(X_{\mathfrak{p}}, Y_{\mathfrak{p}}) \\
&= \dim R/\mathfrak{p} - \inf X_{\mathfrak{p}}^* \otimes_{R_{\mathfrak{p}}}^{\mathbf{L}} Y_{\mathfrak{p}},
\end{aligned}$$

where $X^* = \mathbf{R}\mathrm{Hom}_R(X, R)$ and the last equality comes from Theorem 2.5. Using Nakayama lemma for complexes, we have that

$$\begin{aligned}
\dim R/\mathfrak{p} - \inf \mathbf{R}\mathrm{Hom}_R(X, Y)_{\mathfrak{p}} &= \dim R/\mathfrak{p} - \inf Y_{\mathfrak{p}} - \inf X_{\mathfrak{p}}^* \\
&\leq \dim Y + \mathrm{proj.dim} X_{\mathfrak{p}} \\
&\leq \dim Y + \mathrm{proj.dim} X.
\end{aligned}$$

(ii). By (i) we have the following inequality

$$\dim \mathbf{R}\mathrm{Hom}_R(X, Y) \leq \dim R/\mathfrak{p} - \inf Y_{\mathfrak{p}} + \mathrm{proj.dim} X$$

for some $\mathfrak{p} \in \mathrm{Supp} X \cap \mathrm{Supp} Y$. We can use Nakayama lemma again to see that

$$\begin{aligned}
\dim \mathbf{R}\mathrm{Hom}_R(X, Y) &\leq \dim R/\mathfrak{p} - \inf (X \otimes_R^{\mathbf{L}} Y)_{\mathfrak{p}} + \inf X_{\mathfrak{p}} + \mathrm{proj.dim} X \\
&\leq \dim (X \otimes_R^{\mathbf{L}} Y) + \mathrm{proj.dim} X + \sup X.
\end{aligned}$$

This proves the right hand side of (ii).

Consider the Intersection Theorem for X^* and Y , we have

$$\dim Y \leq \dim (X^* \otimes_R^{\mathbf{L}} Y) + \mathrm{proj.dim} X^*.$$

Now, by using Theorem 2.5, we get

$$\dim Y \leq \dim \mathbf{R}\mathrm{Hom}_R(X, Y) - \inf X.$$

Now the assertion holds. □

Corollary 2.7 Let M be a finite R -module with finite projective dimension, and $Y \in \mathcal{D}_b^f(R)$. Then

$$\dim Y \leq \dim \mathbf{R}\mathrm{Hom}_R(M, Y) \leq \dim (M \otimes_R^{\mathbf{L}} Y) + \mathrm{proj.dim} M.$$

3. Cohomological dimension

In [DY], the authors investigate the invariants $-\inf \mathbf{R}\Gamma_{\mathfrak{a}}(X)$ and $\mathrm{cd}(\mathfrak{a}, X)$, where $\mathfrak{a}$ is an ideal of R , $X \in \mathcal{D}_+(R)$. The main purpose of this section is to study $-\inf \mathbf{R}\Gamma_{\mathfrak{a}}(\mathbf{R}\mathrm{Hom}_R(X, Y))$ for $X, Y \in \mathcal{D}_b^f(R)$. Mainly we seek some results consistent with those in section 2. First we recall the following Theorem, cf. [DY; Theorem 3.2 and Proposition 2.5].

Theorem 3.1 If $X \in \mathcal{D}_+(R)$ is not homologically trivial, then

$$-\inf \mathbf{R}\Gamma_{\mathfrak{a}}(X) \leq \mathrm{cd}(\mathfrak{a}, X).$$

Moreover, if $X \in \mathcal{D}_+^f(R)$, then

$$\begin{aligned} -\inf \mathbf{R}\Gamma_{\mathfrak{a}}(X) &= \mathrm{cd}(\mathfrak{a}, X) \\ &= \sup\{\mathrm{cd}(\mathfrak{a}, R/\mathfrak{p}) - \inf X_{\mathfrak{p}} \mid \mathfrak{p} \in \mathrm{Spec}(R)\}. \end{aligned}$$

We are now ready to give a result which is consistent with Theorem 2.6.

Theorem 3.2 Assume $X, Y \in \mathcal{D}(R)$ and that $\mathbf{R}\mathrm{Hom}_R(X, Y)$ is not homologically trivial. The following statements hold.

(i) If $X \in \mathcal{D}_+^f(R)$ with $\mathrm{proj.dim} X < \infty$ and $Y \in \mathcal{D}_b(R)$, then

$$-\inf \mathbf{R}\Gamma_{\mathfrak{a}}(\mathbf{R}\mathrm{Hom}_R(X, Y)) \leq -\inf \mathbf{R}\Gamma_{\mathfrak{a}}(Y) + \mathrm{proj.dim} X;$$

(ii) If $X, Y \in \mathcal{D}_b^f(R)$ with $\mathrm{proj.dim} X < \infty$, then

$$-\inf \mathbf{R}\Gamma_{\mathfrak{a}}(\mathbf{R}\mathrm{Hom}_R(X, Y)) \leq -\inf \mathbf{R}\Gamma_{\mathfrak{a}}(X \otimes_R^{\mathbf{L}} Y) + \mathrm{proj.dim} X + \sup X.$$

Proof. (i). We have

$$\begin{aligned} -\inf \mathbf{R}\Gamma_{\mathfrak{a}}(\mathbf{R}\mathrm{Hom}_R(X, Y)) &= -\inf \mathbf{R}\mathrm{Hom}_R(X, \mathbf{R}\Gamma_{\mathfrak{a}}(Y)) \\ &= -\inf \mathbf{R}\mathrm{Hom}_R(X, R \otimes_R^{\mathbf{L}} \mathbf{R}\Gamma_{\mathfrak{a}}(Y)) \\ &= -\inf \mathbf{R}\mathrm{Hom}_R(X, R) \otimes_R^{\mathbf{L}} \mathbf{R}\Gamma_{\mathfrak{a}}(Y) \\ &\leq -\inf X^* - \inf \mathbf{R}\Gamma_{\mathfrak{a}}(Y), \end{aligned}$$

where the first equality holds by [Y1; Proposition 2.3] and the third one is Tensor-Hom evaluation. Now the proof of (i) is finished.

(ii). Assume $X, Y \in \mathcal{D}_b^f(R)$, so that $\mathbf{R}\mathrm{Hom}_R(X, Y) \in \mathcal{D}_+^f(R)$ and thus, by Theorem 3.1 and the proof of Theorem 2.6(i),

$$-\inf \mathbf{R}\Gamma_{\mathfrak{a}}(\mathbf{R}\mathrm{Hom}_R(X, Y)) = \mathrm{cd}(\mathfrak{a}, R/\mathfrak{p}) - \inf X_{\mathfrak{p}}^* - \inf Y_{\mathfrak{p}}$$

for some $\mathfrak{p} \in \mathrm{Supp} X \cap \mathrm{Supp} Y$. Now with the same argument as in Theorem 2.6(ii) and using Theorem 3.1 again, we get the result. $\square$

To prove a similar statement as left side equality of Theorem 2.6(ii), one need to show the Intersection Theorem for local cohomology. That is if $X, Y \in \mathcal{D}_b^f(R)$ and $\mathrm{proj.dim} X < \infty$, then

$$-\inf \mathbf{R}\Gamma_{\mathfrak{a}}(Y) \leq -\inf \mathbf{R}\Gamma_{\mathfrak{a}}(X \otimes_R^{\mathbf{L}} Y) + \mathrm{proj.dim} X.$$

Although the proof is not known in general, for the authors, it may worth noting that there is a proof in case $\mathrm{Supp} Y \subseteq \mathrm{Supp} X$. To proceed, it is easy to show that if $X \in \mathcal{D}_b^f(R)$ with finite projective dimension, then $\mathrm{Supp} X^* \subseteq \mathrm{Supp} X$. Thus $\mathrm{Supp} X = \mathrm{Supp} X^{**} \subseteq \mathrm{Supp} X^* \subseteq \mathrm{Supp} X$ gives $\mathrm{Supp} X^* = \mathrm{Supp} X$. Now, we can state the following result.

Theorem 3.3 Assume $X, Y \in \mathcal{D}_b^f(R)$ with $\mathrm{proj.dim} X < \infty$ and $\mathrm{Supp} Y \subseteq \mathrm{Supp} X$. Then

- (i) $-\inf \mathbf{R}\Gamma_{\mathfrak{a}}(Y) \leq -\inf \mathbf{R}\Gamma_{\mathfrak{a}}(X \otimes_R^{\mathbf{L}} Y) + \mathrm{proj.dim} X.$
- (ii) $-\inf \mathbf{R}\Gamma_{\mathfrak{a}}(Y) + \inf X \leq -\inf \mathbf{R}\Gamma_{\mathfrak{a}}(\mathbf{R}\mathrm{Hom}_R(X, Y)).$

Proof. (i). By using Theorem 3.1, there exists $\mathfrak{p} \in \mathrm{Supp} Y$ such that

$$-\inf \mathbf{R}\Gamma_{\mathfrak{a}}(Y) = \mathrm{cd}(\mathfrak{a}, R/\mathfrak{p}) - \inf Y_{\mathfrak{p}}.$$

As $\mathrm{Supp} Y \subseteq \mathrm{Supp} X$, then

$$\begin{aligned} -\inf \mathbf{R}\Gamma_{\mathfrak{a}}(Y) &= \mathrm{cd}(\mathfrak{a}, R/\mathfrak{p}) - \inf Y_{\mathfrak{p}} \\ &= \mathrm{cd}(\mathfrak{a}, R/\mathfrak{p}) - \inf (X \otimes_R^{\mathbf{L}} Y)_{\mathfrak{p}} + \inf X_{\mathfrak{p}} \\ &\leq -\inf \mathbf{R}\Gamma_{\mathfrak{a}}(X \otimes_R^{\mathbf{L}} Y) + \mathrm{proj.dim} X. \end{aligned}$$

The proof of (i) is finished.

(ii) As $\text{Supp } Y \subseteq \text{Supp } X^\star$, we have

$$\begin{aligned} -\inf \mathbf{R}\Gamma_{\mathfrak{a}}(Y) &\leq -\inf \mathbf{R}\Gamma_{\mathfrak{a}}(X^\star \otimes_R^{\mathbf{L}} Y) + \text{proj.dim } X^\star \\ &= -\inf \mathbf{R}\Gamma_{\mathfrak{a}}(\mathbf{R}\text{Hom}_R(X, Y)) - \inf X, \end{aligned}$$

where the equality holds by Theorem 2.5(a) and so the assertion holds. $\square$

Now, we are able to bring the following result.

Corollary 3.4 Assume that M is a finite R -module with finite projective dimension. Then, for any $Y \in \mathcal{D}_b^f(R)$ with $\text{Supp } Y \subseteq \text{Supp } M$,

$$-\inf \mathbf{R}\Gamma_{\mathfrak{a}}(Y) \leq -\inf \mathbf{R}\Gamma_{\mathfrak{a}}(\mathbf{R}\text{Hom}_R(M, Y)).$$

In particular, if $\text{Supp } M = \text{Spec } R$, then

$$-\inf \mathbf{R}\Gamma_{\mathfrak{a}}(R) \leq -\inf \mathbf{R}\Gamma_{\mathfrak{a}}(\mathbf{R}\text{Hom}_R(M, R)).$$

Acknowledgments. The authors would like to thank the referee for his/her substantial comments.

References

- [A] M. Auslander, *Anneaux de Gorenstein, et torsion en algèbre commutative*, (French) Séminaire d'Algèbre Commutative dirigé par Pierre Samuel, 1966/67. Texte rédigé, d'après des exposés de Maurice Auslander, Marquerite Mangeney, Christian Peskine et Lucien Szpiro. École Normale Supérieure de Jeunes Filles Secrétariat mathématique, Paris 1967.
- [AB] M. Auslander; M. Bridger, *Stable module theory*, Memoirs of the American Mathematical Society, No. 94 American Mathematical Society, Providence, R.I. 1969.
- [BH] W. Bruns; J. Herzog, *Cohen-Macaulay rings*, Cambridge Studies in Advanced Mathematics, **39**. Cambridge University Press, Cambridge, 1993.

- [C] L. W. Christensen, *Gorenstein dimensions*, Lecture Notes in Mathematics, **1747** Springer-Verlag, Berlin, 2000.
- [DY] M. T. Dibaei; S. Yassemi, *Cohomological dimension of complexes*, to appear in Comm. Algebra.
- [Fa] G. Faltings, *Über lokale Kohomologiegruppen hoher Ordnung*, J. Reine Angew. Math. **313** (1980), 43–51.
- [Fo] H.-B. Foxby, *Hyperhomological algebra and commutative algebra*, notes in preparation.
- [H] M. Hochster, *Topics in the homological theory of modules over commutative rings*, Regional Conference Series in Mathematics, No. 24. American Mathematical Society 1975.
- [HL] C. Huneke; G. Lyubeznik, *On the vanishing of local cohomology modules*, Invent. Math. **102** (1990), 73–93.
- [HZ] J. Herzog; N. Zamani, *Duality and vanishing of generalized local cohomology*, preprint 2003.
- [PS] C. Peskine; L. Szpiro, *Dimension projective finie et cohomologie locale*, Inst. Hautes études Sci. Publ. Math. No. 42 (1973), 47–119.
- [R] P. Roberts *Le théorème d’intersection*, C. R. Acad. Sci. Paris Sér. I Math. 304 (1987), 177–180.
- [Y1] S. Yassemi, *Generalized section functors*, J. Pure Appl. Algebra **95** (1994), 103–119.
- [Y2] S. Yassemi, *G-dimension*, Math. Scand. **77** (1995), 161–174.